

Table 4. Five-year average rates of confirmed human-caused mortality and serious injury (SI) involving baleen whale stocks along the Gulf of Mexico Coast, US East Coast, and Atlantic Canadian Provinces, 2006-2010 (number of events attributed to entanglements or vessel collisions by year in parentheses).

Stock	Mean annual mortality and SI rate (PBR ¹ for reference)	Entanglements			Vessel Collisions		
		Annual rate (US waters / Canadian waters)	Confirmed mortalities (2006, 2007, 2008, 2009, 2010)	Confirmed SIs (2006, 2007, 2008, 2009, 2010)	Annual rate (US waters / Canadian waters)	Confirmed mortalities (2006, 2007, 2008, 2009, 2010)	Confirmed SIs (2006, 2007, 2008, 2009, 2010)
Western North Atlantic right whale (<i>Eubalaena glacialis</i>)	3.0 (0.9)	1.8 (1.6 / 0.2)	(1, 1, 0, 0, 2)	(0, 0, 1, 2, 2)	1.2 (0.8 / 0.4)	(4, 0, 0, 0, 1)	(1, 0, 0, 0, 0)
Gulf of Maine humpback whale (<i>Megaptera novaeangliae</i>)	7.8 (2.7)	5.8 (5.2 / 0.6)	(1, 1, 2, 2, 3)	(4, 2, 4, 3, 7)	2.0 (2.0 / 0)	(3, 3, 1, 0, 3)	0
Western North Atlantic fin whale (<i>Balaenoptera physalus</i>)	2.0 (5.6)	0.8 (0.6 / 0.2)	(0, 2, 0, 0, 0)	(1, 1, 0, 0, 0)	1.2 (1.2 / 0)	(0, 2, 1, 1, 2)	0
Nova Scotian sei whale (<i>B. borealis</i>)	1.2 (0.6)	0.6 (0.4 / 0.2)	(0, 0, 1, 0, 0)	(1, 0, 1, 0, 0)	0.6 (0.6 / 0)	(1, 1, 0, 1, 0)	0
Western North Atlantic blue whale (<i>B. musculus</i>)	0 (0.9)	0	0	0	0	0	0
Canadian East Coast minke whale (<i>B. acutorostrata</i>)	2.6 (162)	2.2 (1.0 / 1.2)	(2, 1, 4, 0, 0)	(0, 1, 0, 2, 1)	0.4 (0.4 / 0)	(0, 0, 0, 1, 1)	0
Northern Gulf of Mexico Bryde's whale (<i>B. edeni</i>)	0.2 (0.1)	0	0	0	0.2 (0.2 / 0)	(0, 0, 0, 1, 0)	0

¹ Potential Biological Removal (PBR) (Waring *et al* 2011)