

Table 58. Maryland Department of Natural Resources Coastal Bays trawl survey: index of summer flounder recruitment at age-0. Geometric mean (re-transformed \ln [number per hectare + 1]).

Year	Geo. mean n/tow	Lower 95% CI	Upper 95% CI
1972	34.351	13.426	87.888
1973	10.321	5.529	19.267
1974	12.311	7.516	20.165
1975	3.606	2.547	5.104
1976	4.207	2.833	6.246
1977	4.337	2.728	6.894
1978	5.731	3.959	8.295
1979	6.715	4.077	11.060
1980	7.395	3.953	13.837
1981	8.849	5.544	14.123
1982	3.408	1.663	6.983
1983	17.699	0.031	10223.618
1984	13.310	7.161	24.738
1985	12.843	7.472	22.076
1986	59.526	21.950	161.427
1987	7.584	3.590	16.018
1988	1.763	1.371	2.267
1989	2.855	2.121	3.843
1990	4.733	3.639	6.156
1991	7.337	5.508	9.772
1992	8.487	6.285	11.461
1993	4.145	3.192	5.383
1994	22.311	16.486	30.194
1995	13.067	9.811	17.404
1996	6.493	4.954	8.509
1997	7.997	5.948	10.752
1998	14.983	11.391	19.708
1999	8.565	6.477	11.326
2000	9.874	7.272	13.407

Table 58, continued. Maryland Department of Natural Resources Coastal Bays trawl survey: index of summer flounder recruitment at age-0. Geometric mean (re-transformed \ln [number per hectare + 1]).

Year	Geo. mean n/tow	Lower 95% CI	Upper 95% CI
2001	13.543	9.945	18.442
2002	5.406	4.136	7.066
2003	8.180	6.064	11.035
2004	6.993	5.230	9.350
2005	2.198	1.783	2.709
2006	9.658	7.263	12.843
2007	15.438	11.588	20.573
2008	12.079	9.214	15.834
2009	17.887	13.129	24.368
2010	6.713	5.170	8.717
2011	4.471	3.444	5.804