

## **KILLER WHALE (*Orcinus orca*): Northern Gulf of Mexico Stock**

### **STOCK DEFINITION AND GEOGRAPHIC RANGE**

The killer whale is distributed worldwide from tropical to polar regions (Leatherwood and Reeves 1983). Sightings of these animals in the northern Gulf of Mexico (i.e., U.S. Gulf of Mexico) during 1921-1995 occurred primarily in oceanic waters ranging from 256 to 2,652 m (averaging 1,242m) in the north-central Gulf of Mexico (O'Sullivan and Mullin 1997). More recent sightings from NMFS vessel surveys have also occurred in oceanic waters of the north-central Gulf (Figure 1). Despite extensive shelf surveys (O'Sullivan and Mullin 1997), no killer whales have been reported on the Gulf of Mexico shelf waters other than those reported in 1921, 1985 and 1987 by Katona *et al.* (1988). Killer whales were seen only in the summer during GulfCet aerial surveys of the northern Gulf of Mexico between 1992 and 1998 (Hansen *et al.* 1996; Mullin and Hoggard 2000), were reported from May through June during vessel surveys (Mullin and Fulling 2004; Maze-Foley and Mullin 2006) and recorded in May, August, September and November by earlier opportunistic ship-based sources (O'Sullivan and Mullin 1997).

Although there are only a few records from Gulf of Mexico waters beyond U.S. boundaries (e.g., Jefferson and Schiro 1997, Ortega Ortiz 2002), killer whales almost certainly occur throughout the oceanic Gulf of Mexico (Jefferson *et al.* 2008), which is also composed of waters belonging to Mexico and Cuba where there is currently little information on cetacean species abundance and distribution. U.S. waters only comprise about 40% of the entire Gulf of Mexico, and 65% of oceanic waters are south of the U.S. Exclusive Economic Zone (EEZ).

Different stocks have been identified in the northeastern Pacific based on morphological, behavioral and genetic characteristics (Bigg *et al.* 1990; Hoelzel 1991). There is no information on stock differentiation for the Atlantic Ocean population, although an analysis of vocalizations of killer whales from Iceland and Norway indicated that whales from these areas may represent different stocks (Moore *et al.* 1988). Thirty-two individuals have been photographically identified to date in the northern Gulf of Mexico, with 6 individuals having been sighted over a 5-year period, and 1 whale resighted over 10 years. Three animals have been sighted over a range of more than 1,100km (O'Sullivan and Mullin 1997). The Gulf of Mexico population is provisionally being considered a separate stock for management purposes, although there is currently no information to differentiate this stock from the Atlantic Ocean stock(s). Additional morphological, genetic and/or behavioral data are needed to provide further information on stock delineation.

### **POPULATION SIZE**

The best abundance estimate available for northern Gulf of Mexico killer whales is 28 (CV=1.02; Table 1). This estimate is from a summer 2009 oceanic survey covering waters from the 200m isobath to the seaward extent of the U.S. EEZ.

### **Earlier abundance estimates**

All estimates of abundance were derived through the application of distance sampling analysis (Buckland *et al.* 2001) and the computer program DISTANCE (Thomas *et al.* 1998) to line-transect survey data collected from ships in the oceanic northern Gulf of Mexico (i.e., 200m isobath to seaward

259




Figure 1. Distribution of killer whale sightings from SEFSC vessel surveys during spring 1996-2001, summer 2003 and spring 2004, and summer 2009. All the on-effort sightings are shown, though not all were used to estimate abundance. Solid lines indicate the 100-m and 1,000-m isobaths and the offshore extent of the U.S. EEZ.

extent of the U.S. EEZ) and are summarized in Appendix IV.

From 1991 through 1994, and from 1996 through 2001 (excluding 1998), annual surveys were conducted during spring along a fixed plankton-sampling trackline. Due to limited survey effort in any given year, the survey effort-weighted estimated average abundance of killer whales for all surveys combined was estimated. For 1991 to 1994, the estimate was 277 (CV=0.42) (Hansen *et al.* 1995), and for 1996 to 2001, 133 (CV=0.49) (Mullin and Fulling 2004; Table 1).

During summer 2003 and spring 2004, surveys dedicated to estimating cetacean abundance were conducted along a grid of uniformly-spaced transect lines from a random start. The abundance estimate for killer whales, pooled from 2003 to 2004, was 49 (CV=0.77) (Mullin 2007; Table 1).

#### Recent survey and abundance estimate

During summer 2009, a line-transect survey dedicated to estimating the abundance of oceanic cetaceans was conducted in the northern Gulf of Mexico. Survey lines were stratified in relation to depth and the location of the Loop Current. The abundance estimate for killer whales in oceanic waters during 2009 was 28 (CV=1.02; Table 1).

Table 1. Summary of abundance estimates for northern Gulf of Mexico killer whales. Month, year and area covered during each abundance survey, and resulting abundance estimate ( $N_{best}$ ) and coefficient of variation (CV).			
Month/Year	Area	$N_{best}$	CV
Apr-Jun 1991-1994	Oceanic waters	277	0.42
Apr-Jun 1996-2001 (excluding 1998)	Oceanic waters	133	0.49
Jun-Aug 2003, Apr-Jun 2004	Oceanic waters	49	0.77
Jun-Aug 2009	Oceanic waters	28	1.02

#### Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normal distributed abundance estimate. This is equivalent to the 20th percentile of the log-normal distributed abundance estimate as specified by Wade and Angliss (1997). The best estimate of abundance for killer whales is 28 (CV=1.02). The minimum population estimate for the northern Gulf of Mexico is 14 killer whales.

#### Current Population Trend

There are insufficient data to determine the population trends for this species. The pooled abundance estimate for 2003-2004 of 49 (CV=0.77) and that for 1996-2001 of 133 (CV=0.49) are not significantly different ( $P>0.05$ ), but due to the precision of the estimates, the power to detect a difference is low. The abundance estimate for 1991-1994 was 277 (CV=0.42). The large relative changes in the total abundances of killer whales are probably due to a number of factors. The killer whale is most certainly a resident species in the Gulf of Mexico but probably occurs in low numbers and the survey effort is not sufficient to estimate the abundance of uncommon or rare species with precision. Also, these temporal abundance estimates are difficult to interpret without a Gulf of Mexico-wide understanding of killer whale abundance. The killer whale, like all the other oceanic cetacean species in the Gulf, is a mobile predator and this stock is most likely a transboundary stock. The oceanography of the Gulf of Mexico is quite dynamic, and the spatial scale of the Gulf is small relative to the ability of most cetacean species to travel. Studies based on abundance and distribution surveys restricted to U.S. waters are unable to detect temporal shifts in distribution beyond U.S. waters that might account for any changes in abundance.

#### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive history (Barlow *et al.* 1995).

#### POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of the minimum population size, one half the maximum net productivity rate and a recovery factor (MMPA Sec. 3.16 U.S.C. 1362; Wade and Angliss 1997). The minimum

population size is 14. The maximum productivity rate is 0.04, the default value for cetaceans. The recovery factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP), is assumed to be 0.5 because the stock is of unknown status. PBR for the northern Gulf of Mexico killer whale is 0.1.

#### **ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY**

There has been no reported fishing-related mortality of a killer whale during 1998-2010 (Yeung 1999; 2001; Garrison 2003; Garrison and Richards 2004; Garrison 2005; Fairfield Walsh and Garrison 2006; Fairfield-Walsh and Garrison 2007; Fairfield and Garrison 2008; Garrison *et al.* 2009; Garrison and Stokes 2010; 2011). However, during 2008 there was 1 killer whale released alive with no serious injury after an entanglement interaction with the pelagic longline fishery (Garrison *et al.* 2009).

#### **Fisheries Information**

The commercial fishery which potentially could interact with this stock in the Gulf of Mexico is the Atlantic Ocean, Caribbean, Gulf of Mexico large pelagic longline fishery (Appendix III). Pelagic swordfish, tunas and billfish are the targets of the longline fishery operating in the northern Gulf of Mexico. There were no reports of mortality or serious injury to killer whales by this fishery during 1998-2010 (Yeung 1999; 2001; Garrison 2003; Garrison and Richards 2004; Garrison 2005; Fairfield Walsh and Garrison 2006; Fairfield-Walsh and Garrison 2007; Fairfield and Garrison 2008; Garrison *et al.* 2009; Garrison and Stokes 2010; 2011). However, on 17 May 2008 there was 1 killer whale released alive with no serious injury after an entanglement interaction with the pelagic longline fishery (Garrison *et al.* 2009). This was the second observed interaction between a killer whale and this fishery and the first observed interaction within the Gulf of Mexico. During 15 April – 15 June 2008 observer coverage in the Gulf of Mexico was greatly enhanced to collect more robust information on the interactions between pelagic longline vessels and spawning bluefin tuna. Resulting observer coverage for this time and area was dramatically higher than typical for previous years (Garrison *et al.* 2009).

#### **Other Mortality**

There were no reported strandings of killer whales in the Gulf of Mexico during 2006–2010 (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 16 November 2011). Stranding data probably underestimate the extent of fishery-related mortality and serious injury because not all of the marine mammals which die or are seriously injured in fishery interactions wash ashore, not all that wash ashore are discovered, reported or investigated, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interactions.

An Unusual Mortality Event (UME) was declared for cetaceans in the northern Gulf of Mexico beginning 1 February 2010; and, as of early 2012, the event is still ongoing. It includes cetaceans that stranded prior to the Deepwater Horizon oil spill (see “Habitat Issues” below), during the spill, and after. During 2010, no animals from this stock were considered to be part of the UME.

#### **HABITAT ISSUES**

The Deepwater Horizon (DWH) MC252 drilling platform, located approximately 50 miles southeast of the Mississippi River Delta in waters about 1500m deep, exploded on 20 April 2010. The rig sank, and for 87 days millions of barrels of oil and gas were discharged from the wellhead until it was capped on 15 July 2010. During the response effort dispersants were applied extensively at the seafloor and at the sea surface (Lehr *et al.* 2010; OSAT 2010). In-situ burning, or controlled burning of oil at the surface, was also used extensively as a response tool (Lehr *et al.* 2010). The oil, dispersant and burn residue compounds present ecological concerns. The magnitude of this oil spill was unprecedented in U.S. history, causing impacts to wildlife, natural habitats and human communities along coastal areas from western Louisiana to the Florida Panhandle (NOAA 2011). It could be years before the entire scope of damage is ascertained (NOAA 2011).

Shortly after the oil spill, the Natural Resource Damage Assessment (NRDA) process was initiated under the Oil Pollution Act of 1990. A variety of NRDA research studies are being conducted to determine potential impacts of the spill on marine mammals. These studies have focused on identifying the type, magnitude, severity, length and impact of oil exposure to oceanic, coastal and estuarine marine mammals. The research is ongoing and likely will continue for some time. For continental shelf and oceanic cetaceans, the NOAA-led efforts include: aerial surveys to document the

distribution, abundance, species and exposure of marine mammals and turtles relative to oil from DWH spill; and ship surveys to evaluate exposure to oil and other chemicals and to assess changes in animal behavior and distribution relative to oil exposure through visual and acoustic surveys, deployment of passive acoustic monitoring systems, collection of tissue samples, and deployment of satellite tags on sperm and Bryde's whales.

Aerial surveys have observed Risso's dolphins, spinner dolphins, pantropical spotted dolphins, striped dolphins, bottlenose dolphins and sperm whales swimming in oil in offshore waters (NOAA 2010a). The effects of oil exposure on marine mammals depend on a number of factors including the type and mixture of chemicals involved, the amount, frequency and duration of exposure, the route of exposure (inhaled, ingested, absorbed, or external) and biomedical risk factors of the particular animal (Geraci 1990; NOAA 2010b). In general, direct external contact with petroleum compounds or dispersants with skin may cause skin irritation, chemical burns and infections. Inhalation of volatile petroleum compounds or dispersants may irritate or injure the respiratory tract, which could lead to pneumonia or inflammation. Ingestion of petroleum compounds may cause injury to the gastrointestinal tract, which could affect an animal's ability to digest or absorb food. Absorption of petroleum compounds or dispersants may damage kidney, liver and brain function in addition to causing immune suppression and anemia. Long term chronic effects such as lowered reproductive success and decreased survival may occur (Geraci 1990; NOAA 2010b).

### **STATUS OF STOCK**

The status of killer whales in the northern Gulf of Mexico, relative to OSP, is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population trends for this species. Total human-caused mortality and serious injury for this stock is not known but none has been documented. There is insufficient information available to determine whether the total fishery-related mortality and serious injury for this stock is insignificant and approaching zero mortality and serious injury rate. This is not a strategic stock because it is assumed that the average annual human-related mortality and serious injury does not exceed PBR.

### **REFERENCES CITED**

- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Bigg, M.A., P.F. Olesiuk, G.M. Ellis, J.K.B. Ford and K.C. Balcomb 1990. Social organization and genealogy of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. Pages 383-405 *in*: P. S. Hammond, S. A. Mizroch and G. P. Donovan, (eds.) Individual recognition of cetaceans: Use of photoidentification and other techniques to estimate population parameters. Rep. Int. Whal. Comm., Cambridge. Special Issue 12.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D.L. Borchers and L. Thomas 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press. 432 pp.
- Fairfield-Walsh, C. and L.P. Garrison 2007. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2006. NOAA Tech. Memo. NOAA NMFS-SEFSC-560. 54 pp.
- Fairfield, C.P. and L.P. Garrison 2008. Estimated bycatch of marine mammals and sea turtles in the US Atlantic pelagic longline fleet during 2007. NOAA Tech. Memo. NOAA NMFS-SEFSC-572. 62 pp.
- Fairfield Walsh, C. and L.P. Garrison 2006. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2005. NOAA Tech. Memo. NOAA NMFS-SEFSC-539. 52 pp.
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. NOAA Tech. Memo. NMFS-SEFSC-515. 52 pp.
- Garrison, L.P. 2005. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2004. NOAA Tech. Memo. NMFS-SEFSC-531. 57 pp.
- Garrison, L.P. and P.M. Richards 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. NOAA Tech. Memo. NMFS-SEFSC-527. 57 pp.
- Garrison, L.P., L. Stokes and C. Fairfield. 2009. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2008. NOAA Tech. Memo. NMFS-SEFSC-591, 63 pp.
- Garrison, L.P. and L. Stokes. 2010. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2009. NOAA Tech. Memo. NMFS-SEFSC-607, 64 pp.
- Garrison, L.P. and L. Stokes. 2011. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2010. NOAA Tech. Memo. NMFS-SEFSC-624, 59 pp.

- Geraci, J.R. 1990. Physiologic and toxic effects on cetaceans. pp. 167-197 In: J. R. Geraci and D. J. St. Aubin (eds.) Sea mammals and oil: Confronting the risks. Academic Press, New York. 259 pp.
- Hansen, L.J., K.D. Mullin, T.A. Jefferson and G.P. Scott 1996. Visual surveys aboard ships and aircraft. Pages 55-132 in: R. W. Davis and G. S. Fargion, (eds.) Distribution and abundance of marine mammals in the north-central and western Gulf of Mexico: Final report. Volume II: Technical report. OCS Study MMS 96- 0027. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. .
- Hansen, L.J., K.D. Mullin and C.L. Roden 1995. Estimates of cetacean abundance in the northern Gulf of Mexico from vessel surveys. Southeast Fisheries Science Center, Miami, FL. Contribution No. MIA-94/95-25. 9 pp.
- Hoelzel, A.R. 1991. Analysis of regional mitochondrial DNA variation in the killer whale; Implications for conservation. Pages 225-233 in: A. R. Hoelzel, (ed.) Genetic ecology of whales and dolphins. Rep. Int. Whal. Comm., Cambridge. Special Issue 13.
- Jefferson, T. A. and A. J. Schiro. 1997. Distribution of cetaceans in the offshore Gulf of Mexico. Mammal Rev. 27(1): 27-50.
- Jefferson, T.A., M.A. Webber and R.L. Pitman. 2008. Marine mammals of the world. Academic Press, London. 573 pp.
- Katona, S.K.J.A.B., P.E. Girton and F. Wenzel 1988. Killer whales (*Orcinus orca*) from the Bay of Fundy to the equator, including the Gulf of Mexico. Rit Fiskideild 11: 205-224.
- Leatherwood, S. and R.R. Reeves 1983. The Sierra Club handbook of whales and dolphins. Sierra Club Books, San Francisco. 302 pp.
- Lehr, B., S. Bristol and A. Possolo, eds. 2010. Oil budget calculator: Deepwater Horizon. Technical documentation. Prepared by the Federal Interagency Solutions Group, Oil Budget Calculator Science and Engineering Team for the National Incident Command. Available from: [http://www.restorethegulf.gov/sites/default/files/documents/pdf/OilBudgetCalc\\_Full\\_HQ-Print\\_111110.pdf](http://www.restorethegulf.gov/sites/default/files/documents/pdf/OilBudgetCalc_Full_HQ-Print_111110.pdf)
- Maze-Foley, K. and K.D. Mullin 2006. Cetaceans of the oceanic northern Gulf of Mexico: Distributions, group sizes and interspecific associations. J. Cetacean Res. Manage. 8(2): 203-213.
- Moore, S.E., J.K. Francine, A.E. Bowles and K.B. Ford 1988. Analysis of calls of killer whales, *Orcinus orca*, from Iceland and Norway. Rit. Fiskideild 11: 225-250.
- Mullin, K.D. 2007. Abundance of cetaceans in the oceanic Gulf of Mexico based on 2003-2004 ship surveys. Available from: NMFS, Southeast Fisheries Science Center, P.O. Drawer 1207, Pascagoula, MS 39568, 26 pp.
- Mullin, K.D. and G.L. Fulling 2004. Abundance of cetaceans in the oceanic northern Gulf of Mexico. Mar. Mamm. Sci. 20(4): 787-807.
- Mullin, K.D. and W. Hoggard 2000. Visual surveys of cetaceans and sea turtles from aircraft and ships. Pages 111-172 in: R. W. Davis, W. E. Evans and B. Würsig, (eds.) Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: Distribution, abundance and habitat associations. Volume II: Technical report. Minerals Management Service, Gulf of Mexico OCS Region, New Orleans. OCS Study MMS 96-0027.
- NOAA. 2010a. Frequently asked questions about marine mammal rescue and intervention plans in response to the Deepwater Horizon oil spill. Available from: [http://sero.nmfs.noaa.gov/sf/deepwater\\_horizon/20100726\\_FINAL\\_FAQDWH\\_NOAA\\_marine\\_mammal\\_intervention\\_and\\_rescue.pdf](http://sero.nmfs.noaa.gov/sf/deepwater_horizon/20100726_FINAL_FAQDWH_NOAA_marine_mammal_intervention_and_rescue.pdf)
- NOAA. 2010b. Effects of oil on marine mammals and sea turtles. Available from: [http://sero.nmfs.noaa.gov/sf/deepwater\\_horizon/Marine\\_mammals\\_turtles\\_FACT\\_SHEET.pdf](http://sero.nmfs.noaa.gov/sf/deepwater_horizon/Marine_mammals_turtles_FACT_SHEET.pdf).
- NOAA. 2011. Public scoping for preparation of a programmatic environmental impact statement for the Deepwater Horizon BP Oil Spill. Available from: <http://www.gulfspillrestoration.noaa.gov/wp-content/uploads/2011/04/Public-DWH-PEIS-Scoping-Review-Document1.pdf>
- Operational Science Advisory Team (OSAT). 2010. Summary report for sub-sea and sub-surface oil and dispersant detection: Sampling and monitoring. Prepared for P. F. Zukunft, RADM, U.S. Coast Guard, Federal On-Scene Coordinator, Deepwater Horizon MC252, December 17, 2010. Available from: [http://www.restorethegulf.gov/sites/default/files/documents/pdf/OSAT\\_Report\\_FINAL\\_17DEC.pdf](http://www.restorethegulf.gov/sites/default/files/documents/pdf/OSAT_Report_FINAL_17DEC.pdf)
- Ortega Ortiz, J. G. 2002. Multiscale analysis of cetacean distribution in the Gulf of Mexico. Ph.D. dissertation. Texas A&M University, College Station. 170 pp.
- O'Sullivan, S. and K.D. Mullin 1997. Killer whales (*Orcinus orca*) in the northern Gulf of Mexico. Mar. Mamm. Sci. 13(1): 141-147.

- Thomas, L., J.L. Laake, J.F. Derry, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, S. Strindberg, S.L. Hedley, F.F.C. Marques, J.H. Pollard and R.M. Fewster 1998. Distance 3.5. Research Unit for Wildlife Population Assessment, University of St. Andrews, St. Andrews, UK.
- Wade, P.R. and R.P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Yeung, C. 1999. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. NOAA Tech. Memo. NMFS-SEFSC-430. 26 pp.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo., NMFS, Southeast Fisheries Science Center, Miami, FL. NMFS-SEFSC-467. 43 pp.