

Table 1. (A) Estimated dead encounters by gear and year, based on Miller and Shepherd (2011), only for FMPs that will be included in NERO's batched consultation; (B) Bycatch estimates, standard errors, and minimum bycatch. Bycatch estimates are normally distributed, with the lower bound truncated by the observed number of discards, labeled as Minimum. Note that Minimum is not a whole number because of partitioning a whole number by the proportion of adults and subadults in each fishery.

A

Year	Estimated Dead Encounters	
	Sink	
	Gillnet	Otter Trawl
2006	234.4	76.5
2007	344.8	70.7
2008	137.3	60.8
2009	319.9	57.9
2010	191.0	69.4
average	247.7	67.1

B

Gear/Year	Adults			Sub Adults		
	Mean	Standard Error	Minimum	Mean	Standard Error	Minimum
<u>Gillnet</u>						
2006	446.0	170.6	33.5	1,166.0	104.8	87.5
2007	613.2	234.6	30.2	1,602.9	144.9	78.8
2008	237.5	90.8	11.9	620.7	87.7	31.1
2009	568.2	217.4	27.9	1,485.2	131.2	73.1
2010	306.6	117.3	13.8	801.4	99.6	36.2
<u>Trawl</u>						
2006	368.4	95.2	5.8	1,425.3	181.4	22.2
2007	338.1	87.4	11.9	1,307.8	162.0	46.1
2008	285.9	73.9	5.8	1,106.1	135.9	22.2
2009	274.9	71.0	9.9	1,063.3	128.5	38.1
2010	322.5	83.4	21.6	1,247.8	161.3	83.4